

Scotland's Rural College

Use of Infra-Red Thermography to non-invasively assess neonatal piglet temperature

Schmitt, Oceane; O'Driscoll, Keelin

Published in:

Translational Animal Science

DOI:

[10.1093/tas/txaa208](https://doi.org/10.1093/tas/txaa208)

Print publication: 01/01/2021

Document Version

Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Schmitt, O., & O'Driscoll, K. (2021). Use of Infra-Red Thermography to non-invasively assess neonatal piglet temperature. *Translational Animal Science*, 5(1), [txaa208]. <https://doi.org/10.1093/tas/txaa208>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Use of Infra-Red Thermography to non-invasively assess neonatal piglet temperature¹

Oceane Schmitt^{*,†,‡} and Keelin O'Driscoll^{*}

^{*} *Pig Development Department, Teagasc Animal and Grassland Research and Innovation Centre, Fermoy, Ireland*

[†] *Department of Animal Production, Easter Bush Veterinary Centre, Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Roslin, Midlothian, UK*

[‡] *Animal Behaviour and Welfare Team, Animal and Veterinary Sciences Research Group, SRUC, West Mains Road, Edinburgh, UK*

¹ **Acknowledgements:** This research was funded by the Irish Department of Agriculture, Food and the Marine through the FIRM/RSF/CoFoRD 2013 Research Call (project no. 13S428). We would like to thank the farm staff of the Moorepark pig research unit for their help during data collection.

© The Author(s) 2020. Published by Oxford University Press on behalf of the American Society of Animal Science.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

ABSTRACT

Hypothermia is risk factor for piglet neonatal mortality, especially for low birth weight piglets. Piglets with intra-uterine growth retardation (IUGR) also have a higher mortality risk at birth. This study aimed to validate Infra-Red Thermography (IRT) as an alternative to rectal temperature (RT) to measure piglet temperature in the hour post-partum, and to identify piglets with thermoregulation difficulties. At birth (6.3 ± 0.35 min post-partum), 67 piglets were dried, weighed, scored for growth retardation (IUGR; 0-3), and isolated in a plastic box where IRT images were taken, followed by RT. Piglets were then returned to the farrowing pen, and the process repeated at 15, 30 and 60 min post-partum. Piglets were ranked according to their weight (quartiles: 0.57-1.27 kg, 1.27–1.5 kg, 1.5-1.74 kg, 1.74-2.44 kg). Temperatures (ear base and tip; minimum, maximum and average of back) were extracted from IRT images (Thermacam Researcher Pro 2.0). Pearson correlations between temperature measures were calculated, and the effect of time, IUGR score, and weight were included in linear mixed models (SAS 9.4). RT was correlated with all IRT data across time points ($P < 0.05$); correlations were strongest with the ear base, and weakest with the ear tip and minimum back temperature. Both IUGR score and weight rank affected ear base ($P < 0.05$) and rectal temperatures ($P < 0.05$). The lightest piglets, and piglets with severe IUGR had the lowest temperature, relative to their counterparts. Indeed, differences between all weights categories were significant for RT. Piglets with the lowest weight (0.27 – 1.27 kg) had lower ear base temperatures than piglets in the third quartile (1.5–1.74 kg; 35.2 ± 0.36 °C vs. 36.5 ± 0.35 °C, $t_{64,9} = -4.51$, $P < 0.001$) and than heaviest piglets (1.74 – 2.44 kg; 35.2 ± 0.36 °C vs. 36.4 ± 0.36 °C, $t_{70,4} = -3.97$, $P < 0.005$). Overall, piglets with severe IUGR (score

3) had a lower RT than normal piglets (score 0; 35.8 ± 0.46 °C vs. 37.2 ± 0.42 °C, $t_{43,1} = 3.16$, $P < 0.05$) and piglets with mild IUGR (score 1; 35.8 ± 0.46 °C vs. 37.1 ± 0.40 °C, $t_{45,3} = 2.92$, $P < 0.05$); and they also had lower temperature at the base of the ear than normal piglets (35.1 ± 0.42 °C vs. 36.3 ± 0.36 °C, $t_{63,1} = 3.01$, $P < 0.05$). These results confirmed that IRT is an interesting non-invasive tool for assessing neonatal piglets' thermoregulatory abilities and could be used in research investigating successful interventions for piglets at risk of hypothermia.

Keywords: Pigs; infra-red thermography, rectal temperature, thermoregulation, growth retardation, birth weight

List of abbreviations used:

IRT: Infra-Red Thermography

IUGR: Intra-Uterine Growth Retardation

RT: Rectal Temperature

Accepted Manuscript

INTRODUCTION

Hypothermia is a significant cause of neonatal mortality in piglets (Muns et al., 2016). Indeed a series of studies which investigated piglet survival found that piglets which died before weaning had lower temperatures during the first 24 h (Baxter et al., 2008; Baxter et al., 2009; Baxter et al., 2012). Low birth weight piglets have difficulties in maintaining their body temperature, and in recovering from the initial drop in body temperature that happens normally during the hour post-partum (Herpin et al., 2004, 2002; Muns et al., 2016). In addition, some piglets do not achieve optimal growth during gestation (intra-uterine growth retardation; IUGR) and have two to four times higher risk of dying before weaning than normal piglets, depending on the severity of their condition (Hansen et al., 2019). These piglets often show a disproportional allometry at birth (i.e. abnormally long and thin body; Baxter et al., 2008; Hales et al., 2013), and a ‘dolphin-shape’ head. Interventions such as drying piglets at birth and placing them near a heat source (heated lamp or mat) help them to increase and maintain their body heat and avoid hypothermia. In addition, providing an energy supplement should enhance the thermal status of piglets (Muns *et al.*, 2010). However, providing supplemental energy to piglets is rather costly and labour intensive. Thus, piglets most in need of additional support around the time of birth should be identified, as well as the time at which the benefit is maximised (e.g. if there is a time when their temperature is likely to drop). Monitoring of piglets’ temperature during the first hour post-partum could help to identify this critical time-point.

The use of a digital thermometer for measuring rectal temperature is considered the gold standard and has been widely used in research on piglet viability. Recently, infra-red thermography (IRT) imaging has gained interest as a non-invasive technique to measure thermal status of piglets (e.g. Kammersgaard et al., 2011; Tabuaciri et al., 2012; Soerensen and Pedersen, 2015; Sasaki et al., 2016). Kammersgaard et al. (2013) validated the use of IRT

for measuring piglets' temperature by taking IRT images and rectal temperatures of piglets at different times between birth and 48 h post-farrowing. Furthermore, Tabuaciri *et al.* (2012) and Kammergaard *et al.* (2013) both found that the temperature of the base of neonatal piglets' ears (or maximum ear temperature) was most strongly correlated with their rectal temperature. The present study investigated the use of an IRT camera to measure piglets' temperature at several time points during the hour after birth, relative to the rectal temperature (gold standard). This study attempted to maximise control on factors of influence (handling, environment, timing, behaviour) of the temperature at the time of image acquisition. In addition, this study compared known characteristics of piglets' mortality, i.e. weight and IUGR, in order to determine if one is more prevailing in the failure to thermoregulate within 1 h post-partum. We hypothesised that the low birth weight piglets and piglets suffering IUGR would have lower temperatures across time compared to heavier and normal piglets.

MATERIAL AND METHODS

Ethical approval

Ethical approval for this study was granted by the Teagasc Animal Ethics Committee (approval no. TAEC162/2017). The experiment was carried out in accordance with Irish legislation (SI no. 543/2012) and the EU Directive 2010/63/EU for animal experimentation.

Animals and management

This study was conducted in July 2018 in the pig research facilities of the Animal and Grassland Research and Innovation Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland. The experiment involved 67 piglets from 8 litters ($n = 8.4 \pm 0.71$ piglets / litter). Sow genetic background was pure Large White and piglet genetic background was Large White x Duroc.

There were no primiparous sow included in the study: four sows were in their second parity, two sows were in their third parity and two sows were in their fifth parity. Four of the litters were “large litters” as they had over 14 piglets born (two had 16 and, two 17 piglets born alive), and the other four litters had between 7 and 12 piglets born alive. Only two litters had all the piglets involved in the study. This was due to the timing of farrowing, as some piglets were born before the researcher entered the farrowing room on the experimental day. Animals were managed in conventional farrowing pens (250 × 181 cm) containing a sow crate (225 × 60 cm), a heat pad (155 × 37 cm; 2/3 covered), and a water cup and a feeder for piglets.

Thermal image capture

At birth, piglets were dried and isolated in a clear plastic storage box (unknown brand, bought at Toss Bryan, Fermoy, Ireland; 55.4 x 38.0 x 31.7 cm) before acquiring the first thermal image, using a FLIR T420 Infra-Red camera (thermal resolution: 320 x 240, measurement accuracy: $\pm 2^{\circ}\text{C}$, thermal sensitivity: $< 0.04^{\circ}\text{C}$; FLIR Systems, Wilsonville, Oregon, USA). They were then immediately weighed, and scored for intra-uterine growth retardation (IUGR; 0-3). This score was attributed based upon on the number of characteristics associated with intra-uterine growth retardation which were displayed by the piglets. These characteristics were classified by Hales et al. (2015) as having a dolphin-shape head, bulging eyes and wrinkles around the snout (0 = absence of characteristic, 1 = presence of the characteristic). An overall score of 0 indicated no IUGR while a score of 3 indicated severe IUGR.

The plastic box represented a controlled environment for taking IRT images, because it prevented air flow over the piglets. Before acquiring pictures, reflected room temperature was measured as the mean temperature of a crunched aluminium foil (emissivity = 1). The

room temperature was recorded using the same room thermometer (LCD type min/max thermometer, Manotherm; sourced from Ark Animal Care, Newbridge, Ireland) at the time of each image acquisition. These measures allowed to confirm that the room temperature was controlled (each room temperature was individually controlled by Big Farm net program; Big Dutchman AG, Vechta, Germany) and that animals did not suffer heat stress. The skin emissivity of the pig was set at 0.98, as validated by Soerensen et al. (2014). These parameters are important for the correct analysis of the thermal images, as they are used by the software to calculate the subject temperature. Thermal images were acquired at birth (6.3 ± 0.35 min post-partum), 15 min, 30 min and 60 min post-partum, always followed by the taking of a rectal temperature. Images of the piglets' backs were taken at 1 m distance from the piglet with an angle of 75° (Supplementary Material, Fig. S1a). The consistency of these parameters was ensured by adopting the exact same position and respecting landmarks placed on the floor when capturing images. Thermal images were always taken before rectal temperature in order to minimise handling of the piglets, and potential transmission of the experimenter's heat. In addition, the experimenter wore plastic gloves to further ensure insulation of her hands' heat and minimise handling bias. The time spent handling the piglet was recorded, especially at birth when the piglets had to be dried. Rectal temperature measurement took less than 1 min to be obtained, using a digital thermometer (VedoFamily, Pic Solution, Italy).

Animal behaviour

The behaviour of each individual piglet was recorded at each time of image acquisition: the behaviours "walking" (i.e. locomotor activity), "suckling" (i.e. active at the udder, with a teat in the mouth), "huddling" (i.e. sleeping or resting in contact with one or more siblings) and "being on the heat pad" (i.e. resting or active in the heat pad area) were scored as present (score 1) or absent (score 0).

Thermal image analysis

Thermal images were processed with Thermacam Researcher Pro 2.0. Emissivity, reflected temperature and room temperature were modified for each image so that calculated temperatures were accurate. Point measurements were placed at the bases and the tips of piglets' ears, and an area was drawn on their back between the shoulders and the rump (Supplementary Material, Fig. S1b). From this area, the minimum, maximum and average back temperatures were recorded. Temperature data were then entered in an Excel file and analysed as normal.

The reliability of the thermal image analysis was assessed with the intraclass correlation coefficients on SPSS Statistics 24 (IBM corp., Armonk, NY, USA). The same experimenter rated several images of the same piglet (inter-image reliability): coefficients were 0.47 for minimum, 0.95 for maximum, 0.96 for average, 0.89 for ear base and 0.82 for ear tip. Then the same experimenter rated the same images several times (intra-observer reliability): coefficients were 0.44 for minimum, 1.00 for maximum, 1.00 for average, 0.99 for ear base and 0.86 for ear tip.

Statistical analysis

Data were analysed using the software SAS 9.4. The experimental unit was the piglet. Significant terms were determined when the p-value was below 0.05, and tendencies were determined when the p-value was between 0.05 and 0.1.

Pearson correlation tests were performed to investigate the relationships between rectal temperature, birth weight and temperatures obtained from the thermal images of the piglets' ears (tip and base) and back (minimum, maximum and average). Correlations were

characterised “very strong” if the coefficient $r > 0.8$, “strong” if $0.8 > r > 0.5$, “moderate” if $0.5 > r > 0.3$ and “weak” if $0.3 > r$.

Whether there was a significant difference in birthweight of piglets with different IUGR scores was tested using a general linear model (GLM, PROC MIXED). The model included fixed effects of IUGR score and sex, and sow as a random effect.

GLMs were also used for the analysis of all temperature data. We initially analysed the effect of IUGR score on temperature measurements. Intra-uterine growth retardation score, time post-partum, the interaction, the sex of the piglets, and whether or not the piglet performed suckling behaviour, huddling, was located on the heat pad, or was active immediately before being removed for image analysis (i.e. running, playing etc.) and their interaction were included as fixed effects. The random effect of sow and the repeated effect of time were taken into account in all models.

Piglets were then ranked into one of 4 quartiles, based on their birth weight. A similar analysis to that carried out for the effect of IUGR score was carried out; instead of categorising the piglets within an IUGR score, they were categorised within a weight category (very light = 0.53 – 1.23 Kg; light = 1.27 – 1.45 Kg; heavy = 1.50 – 1.72 Kg; very heavy = 1.74 – 2.44 Kg).

RESULTS

Relationship between IUGR score and weight

There was an effect of IUGR score on piglet birth-weight ($P < 0.001$; Table 1). The range of weights overlapped somewhat, and there was no difference in the weight of piglets with IUGR score of 0 or 1. Nevertheless, there was a significant difference between piglets with an IUGR score of 0 and 2 ($t_{60.4} = 3.38$, $P < 0.01$) or 3 ($t_{60.5} = 4.97$, $P < 0.001$), and between piglets with a score of 1 or 3 ($t_{62.7} = 4.23$, $P < 0.001$). Piglets with a score of 1 tended to be heavier than those with a score of 2 ($t_{62.8} = 2.51$, $P = 0.07$), and of score 2 tended to be

heavier than those of score 3 ($t_{60.4} = 2.39$, $P = 0.09$). Therefore, the investigation of the effect of IUGR was quite different from the investigation of the effect of birth weight (category thresholds based on quartiles of the variable weight).

Correlations between thermal data

The correlations between rectal temperature, and birth weight and the thermal data acquired from the IRT images, at each different time of data collection (i.e. birth, 15 min, 30 min, and 60 min post-partum) are presented in Table 2. Rectal temperature was positively correlated with all IRT temperatures at all times ($P < 0.01$). Moderate correlations coefficients (0.3 to 0.6) were found for the minimum temperature of the back and for the temperature of the ear tip; and strong correlations coefficients (0.5 to 0.9) were found for the temperature of the ear base and for the maximum and average temperatures of the back.

Birth weight was positively and moderately correlated to rectal temperature at all times (Table 2). It was also moderately correlated with ear tip and minimum back temperature at birth, and with ear base and maximum back temperature at 30 min and 60 min post-partum. Therefore, the effect of weight on temperature data was studied in more details by using weight ranks (based on weight quartiles) in the analyses.

Effect of time

The effects of time were the same for both analyses of the effect of IUGR score, and weight rank (see Table 3 for the analysis on weight rank, and Table 4 for the analysis on IUGR score). The rectal temperature dropped between birth and 15 min post-partum ($P < 0.001$) and increased again between 30 min and 60 min post-partum ($P < 0.001$). The same pattern was observed with the temperature at the base of the ear (birth vs. 15 min: $P < P < 0.001$; 30 min vs. 60 min: $P < 0.001$) and the tip of the ear (birth vs. 15 min: $P < 0.001$; 30

min vs. 60 min: $P < 0.01$). However, the temperature of the ear tip started to rise again between 15 min and 30 min post-partum ($P < 0.001$).

The minimum temperature of the back also decreased between birth and 15 min post-partum ($P < 0.005$), and increased between 15 min and 60 min post-partum ($P < 0.005$). However, the maximum and average temperatures of the back increased constantly across the hour post-partum. Differences were significant between every time points for the maximum temperature of the back (birth vs. 15 min post-partum: $P < 0.005$; 15 min vs. 30 min post-partum: $P < 0.001$; 30 min vs. 60 min post-partum: $P < 0.001$).

Effect of birth weight category

As hypothesised, there was an effect of birth weight rank on rectal temperature ($F_{3, 44.2} = 14.03$; $P < 0.001$; Table 3). Indeed, there were differences between all rank pairs ($P < 0.05$), except between the two heaviest ranks (Table 3). A similar effect of birth weight rank was found on the temperature of the base of the ear ($F_{3, 65.2} = 8.5$, $P < 0.001$; Table 3). Indeed, piglets in the first quartile (lowest weight; 0.27 – 1.27 kg) had lower ear base temperatures than piglets in the third quartile (1.5 – 1.74 kg) (35.2 ± 0.36 °C vs. 36.5 ± 0.35 °C, $t_{64.9} = -4.51$, $P < 0.001$) and piglets in the fourth quartile (1.74 – 2.44 kg) (35.2 ± 0.36 °C vs. 36.4 ± 0.36 °C, $t_{70.4} = -3.97$, $P < 0.005$), and piglets in the second quartile (1.27 – 1.5 kg) had lower ear base temperature than piglets in the third quartile (35.27 ± 0.35 °C vs. 36.5 ± 0.35 °C, $t_{62.6} = -2.78$, $P < 0.05$). There was an overall effect of rank for the average ($F_{3, 59} = 4.27$; $P < 0.01$) or maximum ($F_{3, 57.4} = 3.62$, $P < 0.05$) temperature of the back, although only piglets in the first quartile had significantly lower temperatures than piglets in the third quartile (maximum: 36.2 ± 0.33 °C vs 36.2 ± 0.33 °C, $t_{57.4}$, $P < 0.05$; average: 33.5 ± 0.38 °C vs. 34.4 ± 0.37 °C, $P < 0.005$). There was no effect of birth weight rank on either the temperature at the ear tip

($F_{3,68.3} = 1.59$; $P = 0.2$), or the minimum temperature recorded on the back ($F_{3,90.9} = 1.46$; $P = 0.2$).

There was an interaction between time and weight rank on the maximum ($F_{9,123} = 2.59$, $P < 0.01$; Fig. 1a) and average ($F_{9,131} = 2.51$, $P < 0.05$; Fig. 1b) temperatures of the back. For both temperatures, the only significant difference was between lightest piglets (first quartile) and piglets in the third weight quartile at 30 min post-partum (maximum: 35.2 ± 0.36 °C vs. 36.4 ± 0.35 °C, $t_{54.3} = -4.34$, $P < 0.005$; average: 33.4 ± 0.4 °C vs. 34.7 ± 0.39 °C, $t_{55.9} = -4.12$, $P < 0.01$).

The interaction between weight rank and time was not significant for rectal temperature ($F_{9,101} = 1.1$, $P = 0.37$), minimum temperature of the back ($F_{9,132} = 0.71$; $P = 0.70$), or temperature at the base ($F_{9,125} = 1.01$, $P = 0.44$) or the tip ($F_{9,134} = 0.97$; $P = 0.47$) of the ear. However, within each time point there were differences in rectal temperature between the weight ranks (birth: $F_{3,42.3} = 7.31$, $P < 0.001$; 15 min post-partum: $F_{3,45.9} = 15.23$, $P < 0.001$; 30 min post-partum: $F_{3,48.5} = 14.6$, $P < 0.001$; 60 min post-partum: $F_{3,45.7} = 5.42$, $P < 0.005$) (Fig. 2). These differences were mainly due to the lightest piglets (first quartile) having lower rectal temperature than heaviest piglets (third and fourth quartiles) at birth (36.6 ± 0.43 °C vs. 38.0 ± 0.40 °C, $t_{41.6} = -4.07$, $P < 0.01$; 36.6 ± 0.43 °C vs. 38.0 ± 0.42 °C, $t_{44.8} = -3.91$, $P < 0.05$; respectively), at 15 min post-partum (35.2 ± 0.40 °C vs. 36.8 ± 0.37 °C, $t_{47.8} = -5.66$, $P < 0.001$; 35.2 ± 0.40 °C vs. 36.9 ± 0.38 °C, $t_{48.5} = -5.84$, $P < 0.001$; respectively) and at 30 min post-partum (34.8 ± 0.43 °C vs. 36.8 ± 0.40 °C, $t_{48.7} = -5.69$, $P < 0.001$; 34.8 ± 0.43 °C vs. 36.8 ± 0.40 °C, $t_{48.2} = -6.04$, $P < 0.001$; respectively) (Fig. 2). At 30 min post-partum, piglets in the first quartile also had a lower rectal temperature than piglets in the second quartile (34.8 ± 0.43 °C vs. 36.1 ± 0.41 °C, $t_{49.4} = -3.57$, $P < 0.05$). Similarly, there were differences between the weight categories in ear base temperatures at 15 min ($F_{3,67.6} = 3.53$; $P < 0.05$), 30 min ($F_{3,55.8} = 5.38$; $P < 0.005$) and 60 min post-partum ($F_{3,57.2} = 6.13$; $P < 0.005$). However,

the differences were only significant between piglets in the first quartile and piglets in the third quartile at 30 min post-partum ($t_{55,4} = -3.61$; $P < 0.05$), and between piglets in the first quartile and piglets in the fourth quartile at 60 min post-partum ($t_{58,5} = -3.87$; $P < 0.05$).

Effect of IUGR score

IUGR score had a significant effect on piglets' rectal temperature ($F_{3,43.6} = 4.07$, $P < 0.05$) and temperature at the base of the ear ($F_{3,62.1} = 3.14$, $P < 0.05$) (Table 4). Indeed, piglets with severe IUGR (score 3) had an overall lower rectal temperature than normal piglets (score 0; 35.8 ± 0.46 °C vs. 37.2 ± 0.42 °C, respectively, $t_{43,1} = 3.16$, $P < 0.05$) and piglets with mild IUGR (score 1; 35.8 ± 0.46 °C vs. 37.1 ± 0.40 °C, respectively, $t_{45,3} = 2.92$, $P < 0.05$). Piglets with severe IUGR also had lower temperature at the base of the ear than normal piglets (35.1 ± 0.42 °C vs. 36.3 ± 0.36 °C, respectively, $t_{63,1} = 3.01$, $P < 0.05$). The effect of IUGR score was not significant on any of the ear tip temperature ($F_{3,68.5} = 0.21$, $P = 0.89$) and the back temperatures (minimum: $F_{3,87.7} = 0.4$, $P = 0.76$; average: $F_{3,56.9} = 1.42$, $P = 0.25$; maximum: $F_{3,54.9} = 1.28$, $P = 0.29$) (Table 4).

There was an interaction between time and IUGR score for rectal temperature ($F_{9,105} = 2.42$, $P < 0.05$; Fig. 3). Indeed, at 30 min post-partum, piglets with severe IUGR (score 3) had lower rectal temperature than normal (IUGR score 0) piglets (35 ± 0.49 °C vs. 36.7 ± 0.45 °C, respectively; $t_{46,6} = 3.52$, $P < 0.05$) and piglets with an IUGR score 1 (35 ± 0.49 °C vs. 36.7 ± 0.42 °C, respectively; $t_{50} = 3.64$, $P < 0.05$). There also was an interaction between time and IUGR score on the minimum ($F_{9,131} = 2$, $P < 0.05$; Fig. 4a) and average ($F_{9,136} = 2.09$, $P < 0.05$; Fig. 4b) temperatures of the back, but pair-wise comparisons were not different. There was no interaction between time and IUGR score for the ear temperatures (ear base: $F_{9,128} = 1.45$, $P = 0.18$; ear tip: $F_{9,134} = 0.52$, $P = 0.86$) or the maximum back temperature ($F_{9,128} = 1.85$, $P = 0.06$).

DISCUSSION

This study aimed at validating the use of infra-red thermography (IRT) to assess the thermal status of piglets across the first hour post-partum, to provide an estimate of their thermoregulatory abilities. Correlations with all temperatures obtained from IRT images with rectal temperature confirmed that IRT can be a valid tool to assess thermal status of piglets around birth. More specifically, images of the ears showed the same thermal patterns as rectal temperature over time. The second aim of the study was to identify whether the level of intra-uterine growth retardation (IUGR) and birth weight could influence thermoregulatory ability; the aim was to determine whether the pattern of temperature change over time was affected by weight and IUGR score. The results showed clearly that both factors did indeed influence thermoregulatory abilities of piglets.

Correlations between rectal temperature and ear (base and tip) and back (minimum, maximum and average temperatures of the back area, from shoulders to rumps) temperatures confirmed earlier work showing that infra-red thermography is a valid tool for assessing piglets temperature (Tabuaciri et al., 2012; Kammergaard et al., 2013; Soerensen and Pedersen, 2015). Furthermore, strong correlations with rectal temperatures at each time point suggested that maximum back and ear base temperatures are best locations for approximating piglet body temperature. The levels of correlation found in the present study are similar to the ones found in other studies (Tabuaciri et al., 2012; Kammergaard et al., 2013). Higher correlations could be expected in the present study as the time of acquisition of images (relative to the piglet's birth) and environmental factors (isolation in a plastic box) were controlled, which was not the case in previous work (Kammergaard et al., 2013; Tabuaciri et al., 2012). Indeed, piglets were handled during image acquisition in the study of Kammergaard et al. (2013), which can cause stress and elevation of body temperature, and

the time post-partum was not accounted for in the study of Tabuaciri et al. (2012). Moreover, there are potential confounding factors not accounted for in these studies, such as environmental temperature, piglets' behaviour (e.g. huddling; Llamas Moya et al., 2006), and presence of birth fluids.

During the first hour post-partum, piglets' rectal temperature, the gold standard in core temperature measurement, decreased between birth and 15 min post-partum, then increased again between 30 min and 60 min post-partum. The ear base, ear tip and the minimum back temperatures followed the same pattern. This shows thermoregulation process, i.e. the change of temperature overtime to reach or maintain thermal homeostasis, and such pattern (decrease shortly after birth and increase afterwards) was previously reported by Herpin et al. (2002). However, the maximum and average back temperatures increased steadily overtime, which confirms the findings of Kammersgard et al. (2013) that the back temperature may not provide an accurate estimate of the core body temperature.

The effect of level of IUGR and weights on temperature data were investigated separately, as they describe different populations of piglets. Indeed, even if lower average weights correspond to a greater severity of IUGR, it is important to make a distinction between piglets born with low birth-weight (also called "small for gestational age") and piglets which experience IUGR, because their survival chance and growth potential might be different (Rutherford et al., 2013). In the present study, both the level of IUGR and weight ranks (based on weight quartiles) affected piglets' rectal and ear base temperatures. However, significant differences between the IUGR levels were only found at 30 min post-partum between severe IUGR piglets (score 3) and normal piglets (score 0), whereas weight ranks differed more across time. The second difference was found in the pattern of rectal temperature, as IUGR level did not affect piglets' rectal temperature at birth but weight rank did. Indeed, piglets with the lowest birth weight (0.57 – 1.27 kg) had a lower rectal

temperature compared to heavier piglets from birth until 30 min post-partum. Moreover, at 60 min post-partum, piglets with a body weight above 1.5 kg had a rectal temperature greater than 37°C whereas the rectal temperature of piglets under 1.27 kg was just above 36°C, showing that the later might have some difficulty in ensuring thermoregulation (i.e. piglets should reach the thermal homeostasis of 39°C within 48 h of age; Herpin et al., 2002). The same difference of temperature at 60 min post-partum was also observed between piglets with severe IUGR, compared to all other levels. Altogether, our results suggest that IUGR and low birth weight are two separated conditions, often affecting the same piglets, but which may have separate influences on piglet thermoregulatory abilities. It is possible that a low birth weight, independent of the level of IUGR, is an unfavourable condition for thermoregulation, due to the greater surface to body mass ratio and consequent greater heat loss (Herpin et al., 2002), and that IUGR level may prevent these small piglets from ensuring their thermoregulation within 1 h post-partum. Therefore, affected piglets may require a greater amount of time to reach thermal comfort, or a greater supply of energy from colostrum. The present results may suggest that piglets with low birth weights rather than IUGR should be targeted by energy supplementation interventions.

In conclusion, this study confirmed that infra-red thermography, and especially images of the base of the ear, is a valid non-invasive tool to assess thermal status of neonatal piglets. Images taken during the first hour post-partum could be used to determine piglets with difficulties in maintaining body temperature under the experimental conditions. Further research work could use infra-red thermography to test the effects of energy supplementation on the thermoregulatory abilities of neonatal piglets, in order to identify the most successful timing of supplementation.

DISCLOSURES

The authors declare no conflict of interest.

LITERATURE CITED

- Baxter, E. M., S. Jarvis, R. B. D'Eath, D. W. Ross, S. K. Robson, M. Farish, I. M. Nevison, A. B. Lawrence, and S. A. Edwards. 2008. Investigating the behavioural and physiological indicators of neonatal survival in pigs. *Theriogenology*. 69:773–783.
doi:10.1016/j.theriogenology.2007.12.007.
- Baxter, E. M., S. Jarvis, J. Palarea-Albaladejo, and S. a. Edwards. 2012. The weaker sex? the propensity for male-biased piglet mortality. *PLoS One*. 7. doi:10.1371/journal.pone.0030318.
- Baxter, E. M., S. Jarvis, L. Sherwood, S. K. Robson, E. Ormandy, M. Farish, K. M. Smurthwaite, R. Roehe, A. B. Lawrence, and S. A. Edwards. 2009. Indicators of piglet survival in an outdoor farrowing system. *Livest. Sci.* 124:266–276.
doi:10.1016/J.LIVSCI.2009.02.008. Available from:
<https://www.sciencedirect.com/science/article/pii/S1871141309000614>
- Hales, J., V. A. Moustsen, A. M. Devreese, M. B. F. Nielsen, and C. F. Hansen. 2015. Comparable farrowing progress in confined and loose housed hyper-prolific sows. *Livest. Sci.* 171:64–72. doi:10.1016/j.livsci.2014.11.009.
- Hales, J., V. A. Moustsen, M. B. F. Nielsen, and C. F. Hansen. 2013. Individual physical characteristics of neonatal piglets affect preweaning survival of piglets born in a noncrated system. *J. Anim. Sci.* 91:4991–5003. doi:10.2527/jas2012-5740.
- Hansen, C. F., J. Hales, C. Amdi, and V. A. Moustsen. 2019. Intrauterine growth-restricted piglets defined by their head shape have impaired survival and growth during the suckling period. *Anim. Prod. Sci.* doi:10.1071/AN17581.
- Herpin, P., M. Damon, and J. Le Dividich. 2002. Development of thermoregulation and

neonatal survival in pigs. *Livest. Prod. Sci.* 78:25–45. doi:10.1016/S0301-6226(02)00183-5.

Kammersgaard, T. S., J. Malmkvist, and L. J. Pedersen. 2013. Infrared thermography--a non-invasive tool to evaluate thermal status of neonatal pigs based on surface temperature.

Animal. 7:2026–34. doi:10.1017/S1751731113001778. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24237678>

Kammersgaard, T. S., L. J. Pedersen, and E. Jørgensen. 2011. Hypothermia in neonatal piglets : interactions and causes of individual differences. *J. Anim. Sci.* 89:2073–2085. doi:10.2527/jas.2010-3022.

Llamas Moya, S., L. A. Boyle, P. B. Lynch, and S. Arkins. 2006. Influence of teeth resection on the skin temperature and acute phase response in newborn piglets. *Anim. Welf.* 15:291–297.

Muns, R., M. Nuntapaitoon, and P. Tummaruk. 2016. Non-infectious causes of pre-weaning mortality in piglets. *Livest. Sci.* 184:46–57. doi:10.1016/j.livsci.2015.11.025. Available from: <http://dx.doi.org/10.1016/j.livsci.2015.11.025>

Muns, R., R. de la Torre, P. S. Agostini, X. Manteca, and J. Gasa. 2010. The effect of colostrum supplementation on piglets' body temperature recovery and lactation performance. *J. Anim. Sci.* 88:306.

Rutherford, K. M. D., E. M. Baxter, R. B. D'Eath, S. P. Turner, G. Arnott, R. Roehe, B. Ask, P. Sandøe, V. A. Moustsen, F. Thorup, S. A. Edwards, P. Berg, and A. B. Lawrence. 2013. The welfare implications of large litter size in the domestic pig I: biological factors. *Anim. Welf.* 22:199–218. doi:10.7120/09627286.22.2.199.

Sasaki, Y., K. Furusho, R. Ushijima, T. Tokunaga, R. Uemura, and M. Sueyoshi. 2016. Body Surface Temperature of Suckling Piglets Measured by Infrared Thermography and Its Association with Body Weight Change. *Japan Agric. Res. Q.* 50:361–368.

Soerensen, D. D., S. Clausen, J. B. Mercer, and L. J. Pedersen. 2014. Determining the

emissivity of pig skin for accurate infrared thermography. *Comput. Electron. Agric.* 109:52–58. doi:10.1016/j.compag.2014.09.003. Available from:

<http://www.sciencedirect.com/science/article/pii/S0168169914002178>

Soerensen, D. D., and L. J. Pedersen. 2015. Infrared skin temperature measurements for monitoring health in pigs: a review. *Acta Vet. Scand.* 57:5. doi:10.1186/s13028-015-0094-2. Available from: <http://www.actavetscand.com/content/57/1/5>

Tabuaciri, P., K. L. Bunter, and H.-U. Graser. 2012. Thermal imaging as a potential tool for identifying piglets at risk. In: *AGBU Pig Genetics Workshop*. Armidale, Australia. p. 23–30.

Accepted Manuscript

Figure 1 Maximum (a) and average (b) back temperatures (LS mean \pm S.E.) of piglets in different weight categories. Different letters indicate differences between weight categories (lowercase: $P < 0.05$, uppercase $P < 0.005$). P-value for the effect of the interaction between time and weight rank was $P < 0.01$ for the maximum back temperature, and $P < 0.05$ for the average back temperature.

Figure 2 Rectal temperature (LS mean \pm S.E.) of piglets in different weight rank across the first hour post-partum (effect time \times weight rank: $P > 0.1$).

Figure 3 Rectal temperature (LS mean \pm S.E.) of piglets with different IUGR score across the first hour post-partum (effect time \times IUGR score: $P < 0.05$). Different letters indicate differences between weight categories ($P < 0.05$).

Figure 4 Minimum (a) and average (b) back temperatures (LS mean \pm S.E.) of piglets with different IUGR score. Different letters indicate differences between weight categories (lowercase: $P < 0.05$, uppercase $P < 0.005$). P-value for the effect of the interaction between time and IUGR score was $P < 0.05$ for both the minimum and average back temperatures.

Table 1 Birth weights of piglets classified into each IUGR category (score 0 = no IUGR to score 3 = severe IUGR).

	N	Minimum	Maximum	LS Mean \pm S.E.
IUGR 0	15	1.03	2.44	1.68 \pm 0.09 ^a
IUGR 1	22	0.98	2.08	1.57 \pm 0.09 ^{ab}
IUGR 2	21	0.78	1.79	1.33 \pm 0.08 ^{bc}
IUGR 3	9	0.53	1.56	1.04 \pm 0.12 ^c

Accepted Manuscript

Table 2 Pearson correlations coefficients characterising relationships between rectal temperature, and birth weight and thermal data. Data from thermal images (temperature of the ear tip, ear base, and minimum, maximum and average back temperature) obtained at birth (6 min) and 15 min, 30 min and 60 min post-partum (pp). *P < 0.05, ** P < 0.01; ***P < 0.001

	Back temperatures			Ear temperatures		Birth weight
	Minimum	Maximum	Average	Tip	Base	
Rectal temperature						
6 min pp. ¹ (birth)	0.48**	0.55***	0.57***	0.59***	0.8***	0.47**
15 min pp.	0.37**	0.45***	0.49***	0.28*	0.64***	0.57***
30 min pp.	0.36**	0.86***	0.78***	0.53***	0.86***	0.56***
60 min pp.	0.59***	0.9***	0.87***	0.39**	0.71***	0.36**
Birth Weight						
6 min pp. (birth)	0.28*	0.04	0.1	0.31*	0.21	
15 min pp.	0.22	-0.03	-0.1	0.04	0.1	
30 min pp.	-0.05	0.28*	0.15	0.03	0.32**	
60 min pp.	0.01	0.28*	0.19	0.18	0.32**	

¹ pp. = post-partum

Table 3 Mean \pm S.E. temperatures of neonatal piglets, according to the time post-partum and their birth weight category. Superscript letters indicate significant differences within each variable (^{a,b} = $P < 0.05$, ^{A,B} = $P < 0.005$).

	Rectal Temperature	Ear base temperature	Ear tip temperature	Minimum back temperature	Maximum back temperature	Average back temperature
Effect of time						
Birth (6 min pp. ¹)	37.4 \pm 0.36 ^A	36.1 \pm 0.33 ^{AC}	29.2 \pm 0.32 ^{AC}	29.3 \pm 0.43 ^A	34.9 \pm 0.32 ^A	33.2 \pm 0.35 ^A
15 min post-partum	36.2 \pm 0.34 ^B	35.4 \pm 0.31 ^B	27.7 \pm 0.32 ^B	28.3 \pm 0.43 ^B	35.4 \pm 0.32 ^B	33.5 \pm 0.35 ^A
30 min post-partum	36.2 \pm 0.36 ^B	35.7 \pm 0.32 ^{AB}	28.8 \pm 0.35 ^A	28.7 \pm 0.42	35.9 \pm 0.31 ^C	34.1 \pm 0.35 ^B
60 min post-partum	37.1 \pm 0.37 ^A	36.6 \pm 0.36 ^C	29.8 \pm 0.37 ^C	29.3 \pm 0.44 ^A	36.7 \pm 0.30 ^D	34.9 \pm 0.35 ^C
<i>P-value</i>	$P < 0.001$	$P < 0.001$	$P < 0.001$	$P < 0.001$	< 0.001	< 0.001
Effect of weight category						
< 1.27 kg	35.7 \pm 0.41 ^{Aa}	35.2 \pm 0.36 ^A	28.4 \pm 0.39	28.7 \pm 0.44	35.4 \pm 0.34 ^a	33.5 \pm 0.38 ^A
1.27 - 1.50 kg	36.5 \pm 0.39 ^b	35.7 \pm 0.35 ^a	29.2 \pm 0.37	29.2 \pm 0.43	35.6 \pm 0.34	33.9 \pm 0.37
1.50 - 1.74 kg	37.3 \pm 0.38 ^{Bc}	36.5 \pm 0.35 ^{Bb}	28.9 \pm 0.37	29.0 \pm 0.43	36.2 \pm 0.33 ^b	34.4 \pm 0.37 ^B
> 1.74 kg	37.4 \pm 0.39 ^{Bc}	36.4 \pm 0.36 ^B	28.9 \pm 0.39	28.8 \pm 0.44	35.7 \pm 0.34	33.8 \pm 0.38
<i>P-value</i>	$P < 0.001$	$P < 0.001$	$P = 0.2$	$P = 0.2$	$P < 0.01$	$P < 0.05$

¹ pp. = post-partum

Table 4 Mean (\pm S.E.) temperatures of neonatal piglets, according to the time post-partum and their IUGR score. Superscript letters indicate significant differences within each variable (^{a,b} = $P < 0.05$, ^{A,B} = $P < 0.005$).

	Rectal temperature	Ear base temperature	Ear tip temperature	Minimum back temperature	Maximum back temperature	Average back temperature
Effect of time						
Birth (6 min pp. ¹)	37.4 \pm 0.35 ^A	36.1 \pm 0.33 ^A	29.3 \pm 0.35 ^{AC}	29.2 \pm 0.47 ^A	35.0 \pm 0.32 ^A	33.2 \pm 0.36 ^A
15 min post-partum	36.2 \pm 0.34 ^B	35.3 \pm 0.30 ^B	27.8 \pm 0.35 ^B	28.2 \pm 0.45 ^B	35.3 \pm 0.32 ^A	33.4 \pm 0.37 ^A
30 min post-partum	36.1 \pm 0.35 ^B	35.6 \pm 0.31 ^{ABa}	28.7 \pm 0.38 ^A	28.8 \pm 0.45	35.8 \pm 0.31 ^B	34.0 \pm 0.36 ^B
60 min post-partum	37.0 \pm 0.37 ^A	36.3 \pm 0.37 ^{Ab}	29.8 \pm 0.39 ^C	29.4 \pm 0.47 ^A	36.6 \pm 0.30 ^C	34.8 \pm 0.37 ^C
<i>P-value</i>	$P < 0.001$	$P < 0.001$	$P < 0.001$	$P < 0.001$	< 0.001	< 0.001
Effect of IUGR score						
IUGR 0	37.2 \pm 0.42 ^a	36.3 \pm 0.36 ^a	29.0 \pm 0.40	28.8 \pm 0.46	35.7 \pm 0.34	33.9 \pm 0.39
IUGR 1	37.1 \pm 0.40 ^a	36.1 \pm 0.34	29.0 \pm 0.38	29.1 \pm 0.45	35.7 \pm 0.33	33.9 \pm 0.38
IUGR 2	36.6 \pm 0.38	35.9 \pm 0.34	28.8 \pm 0.38	28.9 \pm 0.45	35.9 \pm 0.33	34.1 \pm 0.38
IUGR 3	35.8 \pm 0.46 ^b	35.1 \pm 0.42 ^b	28.8 \pm 0.49	28.9 \pm 0.50	35.3 \pm 0.39	33.5 \pm 0.44
<i>P-value</i>	$P < 0.05$	$P < 0.05$	$P = 0.89$	$P = 0.76$	$P = 0.29$	$P = 0.25$

¹ pp. = post-partum

Figure 1






Figure 2



Accepted

Figure 3



Accepted

Figure 4

